

# Chapter 3

## Cosmology

### 3.1 Expanding universe

The observable universe is observed to be spatially isotropic, homogeneous and flat. Decomposing the metric as in Eq. (2.2.9), spatial isotropy implies  $B_{\mathbf{a}} = 0$ , spatial homogeneity implies  $A = A(t)$  and  $h_{\mathbf{ab}} = h_{\mathbf{ab}}(t)$  and hence we can redefine the time coordinate such that  $A = 1$ , and spatial isotropy, homogeneity and flatness imply  $h_{\mathbf{ab}} \propto \delta_{\mathbf{ab}}$  where  $\delta_{\mathbf{ab}}$  is the metric on three dimensional Euclidean space. Thus the metric for a spatially isotropic, homogeneous and flat universe takes the form

$$g_{\mathbf{ab}} = e_{\mathbf{a}}^t e_{\mathbf{b}}^t - a^2(t) \delta_{\mathbf{ab}} \quad (3.1.1)$$

or equivalently

$$d\tau^2 = g_{\mathbf{ab}} dx^{\mathbf{a}} dx^{\mathbf{b}} = dt^2 - a^2(t) \delta_{\mathbf{ab}} dx^{\mathbf{a}} dx^{\mathbf{b}} \quad (3.1.2)$$

where  $a(t)$  is the **scale factor** and  $\delta_{\mathbf{ab}}$  is the **comoving** metric. Similarly, the stress tensor takes the form

$$T_{\mathbf{ab}} = \rho(t) e_{\mathbf{a}}^t e_{\mathbf{b}}^t + p(t) (e_{\mathbf{a}}^t e_{\mathbf{b}}^t - g_{\mathbf{ab}}) \quad (3.1.3)$$

Substituting Eqs. (3.1.1) and (3.1.3) into the Einstein equation, Eq. (2.4.9), gives the **Friedmann equations**

$$3 \frac{\dot{a}^2}{a^2} = \rho \quad (3.1.4)$$

$$\frac{\ddot{a}}{a} = -\frac{1}{6} (\rho + 3p) \quad (3.1.5)$$

Restoring  $c$  and  $8\pi G$ , Eq. (3.1.5) can be rewritten in Newtonian style as

$$\ddot{a} = -\frac{G(\rho + 3p) \left(\frac{4}{3}\pi a^3\right)}{c^2 a^2} \quad (3.1.6)$$

with the relativistic source  $(\rho + 3p)/c^2$  reducing to the mass density in the Newtonian limit.

Eq. (3.1.4) relates the curvature of spacetime, in the form of the expansion rate or **Hubble parameter**

$$H \equiv \frac{\dot{a}}{a} \quad (3.1.7)$$

to the energy density. Reexpressing Eqs. (3.1.4) and (3.1.5) in terms of the Hubble parameter gives

$$3H^2 = \rho \quad (3.1.8)$$

$$\dot{H} = -\frac{1}{2}(\rho + p) \quad (3.1.9)$$

and taking the derivative gives

$$\dot{\rho} = -3H(\rho + p) \quad (3.1.10)$$

which can also be derived directly from  $\nabla_{\mathbf{b}}T_{\mathbf{a}}^{\mathbf{b}} = 0$ .

From Eq. (3.1.2), physical spatial distances are given by

$$s = ax \quad (3.1.11)$$

where  $x$  is the comoving coordinate distance. Therefore, for objects moving with the expansion of the universe, i.e. with constant  $x$ , we have

$$\frac{ds}{dt} = \dot{a}x = \frac{\dot{a}}{a}s = Hs \quad (3.1.12)$$

which is **Hubble's law**: comoving objects are receding from us at a speed proportional to their distance from us. Further, at distances  $s > 1/H$  we have  $ds/dt > 1$  and so comoving objects are moving away from us at greater than the speed of light and so are out of causal contact and are said to be beyond the **horizon**. The **observable universe** corresponds to  $s < 1/H$ .