

Homework 13 - Symmetry

- Q13.1. (a) Express $\mathcal{L}_u v^{\mathbf{a}}$ in a coordinate basis and deduce that it is independent of the metric.
 (b) Use Eq. (2.3.20) to show that

$$\mathcal{L}_u \omega_{\mathbf{a}} = u^{\mathbf{b}} \nabla_{\mathbf{b}} \omega_{\mathbf{a}} + \omega_{\mathbf{b}} \nabla_{\mathbf{a}} u^{\mathbf{b}} \quad (\text{Q13.1.1})$$

and check that this is consistent with Eq. (1.2.6).

- Q13.2. (a) Derive Eq. (2.3.21).
 (b) Show that a coordinate basis vector $e_{\alpha}^{\mathbf{a}}$ is a Killing vector if and only if

$$\nabla_{\alpha} g_{\beta\gamma} = 0 \quad (\text{Q13.2.1})$$

for all β, γ , and explain the difference between $\nabla_{\alpha} g_{\beta\gamma}$ and $\nabla_{\mathbf{a}} g_{\mathbf{bc}}$.

- (c) Show that a particle with momentum

$$p_{\mathbf{a}} = m g_{\mathbf{ab}} \frac{dx^{\mathbf{b}}}{dt} \quad (\text{Q13.2.2})$$

and moving freely in a space with Killing vector field $\xi^{\mathbf{a}}$ has conserved quantity $\xi^{\mathbf{a}} p_{\mathbf{a}}$.